

MASSACHUSETTS INSTITUTE OF TECHNOLOGY
ARTIFICIAL INTELLIGENCE LABORATORY

A.I. memo No. 591

April, 1980

INTERFACING THE ONE-DIMENSIONAL SCANNING OF AN IMAGE
WITH THE APPLICATIONS OF TWO-DIMENSIONAL OPERATORS

S. Ullman

Abstract: To interface between the one-dimensional scanning of an image, and the application of a two-dimensional operator, an intermediate storage is required. For a square image of size n^2 , and a square operator of size m^2 , the minimum intermediate storage is shown to be $n \cdot (m - 1)$. An interface of this size can be conveniently realized by using a serpentine delay line. New kinds of imagers would be required to reduce the size of the intermediate storage below $n \cdot (m - 1)$.

This report describes research done at the Artificial Intelligence Laboratory of the Massachusetts Institute of Technology. Support for the laboratory's artificial intelligence research is provided in part by the Advanced Research Project Agency of the Department of Defence under Office of Naval Research contract N00014-75-C-0643 and in part by National Science Foundation Grant MCS77-07569

1. Introduction

Image processing often requires the application of a two-dimensional operator over the image. Imaging devices usually scan the image along a one-dimensional path, and produce a stream of data points. In such cases, an intermediate storage is required, to interface between the one-dimensional scanning and the application of the two-dimensional operator.

In this paper, the interface problem will be examined. In particular, we shall determine the minimum size required of the intermediate storage. For a square image of size n^2 , and a square operator of size m^2 , the minimum intermediate storage is shown to be $n \cdot (m - 1)$. For small-size operators the intermediate storage may not add substantially to the overall size of the system. For example, a recent two-dimensional CCD scanner developed by Texas Instruments (Hall & Awtrey, 1979) incorporates the intermediate storage required for a $3 * 3$ operator on the scanner chip itself. For larger operators, however, the size of the intermediate storage becomes significant. For example, a recent CCD convolution chip constructed by Hughes is designed to support a $26 * 26$ operator (Nudd *et al.*, 1979). It is intended to be applied to images of 1000 points on a side, in which case an intermediate storage of 25000 image points would be required. In such cases the size of the system is determined almost entirely by the intermediate storage requirement. Given the theoretical lower bound on the size of this storage, it is suggested that new kinds of imagers would be required for systems for which size is a dominant consideration.

2. Using a serpentine delay line of size $n \cdot (m - 1)$

A simple method for interfacing a one-dimensional scanner to a two-dimensional operator is shown in Figure 1. The camera c scans the image (assumed to be a square of n^2 points) row by row. The intensity values measured by the camera are fed into a serpentine delay line (S in Figure 1), consisting of $m - 1$ rows, each of length n . The serpentine delay line is tapped in $m - 1$ positions that are connected to the two-dimensional operator M (of size m^2). In addition, there is a direct line from the camera to M , so that there are a total of m

input lines to M.

After the first $m - 1$ rows have been scanned, they fill the intermediate storage S, and the first column of these rows resides in the rightmost column of S, ready to be delivered to M. The scanner then reads point $(m,1)$ in the image, and the first column of m points (point $(1,1)$ to $(m,1)$) is transferred to M. Point $(m,1)$ also enters the delay line, shifting each point in S by one position. The point in the lower-right corner of S is dropped from the memory and lost. Whenever a new point is now read by the scanner, a column of m points is transferred to M. After having scanned m points from the m^{th} row, the entire upper-left square of m^2 points resides in M, and the operator is applied to it. When the scanning of the entire image is completed, the operator has been applied to all the squares of size m^2 in the image.

The intermediate storage S in this scheme is required to hold $n \cdot (m - 1)$ image points. The next section will establish that this requirement cannot be reduced. Even if one uses arbitrary scanning pattern (e.g., the image may be divided into sub-blocks, and each block scanned and operated upon separately), and arbitrary interconnections between S and M (i.e., M may have access to all the points in S), the minimal intermediate storage would still be $n \cdot (m - 1)$.




Figure 1: A serpentine delay line **S**, with $m - 1$ rows of length n each, interfaces a scanner **C** with the operator **M** of size m^2 .

3. The minimal intermediate storage is $n \cdot (m - 1)$

Let the size of the intermediate storage be S . In this part we wish to show that $S \geq n \cdot (m - 1)$.

Notation: We shall think of the image as a grid of $n * n$ squares, $n \geq 2$. Each square in the grid is also called a *point* in the image. A two-dimensional operator of size $m * m$ is a function of m^2 arguments. The function is to be applied to every square of size m^2 points in the image. We shall assume that the size of the operator (m) is at most $\frac{n}{2}$. Finally, for a set P of points, $\|P\|$ will denote the number of points in P .

Definitions: Two distinct points in the image are called *partners* if they can be covered simultaneously by a square of size m^2 . That is, their distance in the image in either the horizontal or the vertical direction is at most m points. Similarly, the partners of a set P of points are the partners of all the points in P , but not including the points in the set P itself. That is,

$$\text{partners}(P) = \bigcup_{i \in P} \text{partners}(i) - P \quad (1)$$

When the scanner has read k points, $k > S$, some of the k points are no longer in the storage, and can no longer be retrieved. Let P be the set of points that have been scanned, but are no longer in the storage. It is not difficult to see that at this stage the partners of the set P must reside in the intermediate storage. The reason is that if points p and q are partners, then there must exist a stage at which both p and q are in the intermediate storage. We can now obtain a lower bound on the size of the intermediate storage by estimating the number of partners of a set of k points. We shall establish the lower bound on the size of the intermediate storage by showing that for a certain choice of k , the number of partners is at least $n \cdot (m - 1)$.

Proposition 1: Let k be the smallest integer such that $k > \frac{n^2}{4}$. For this choice of k , any set of k points has at least $n(m - 1)$ partners.

Proof: By induction on m , the size of the two-dimensional operator. For $m = 1$ the claim holds, since the number of partners is zero. To prove the induction step we shall show that, in an image of size n^2 , when the size of the operator is increased from $m = M$ to $m = M + 1$, (assuming $M + 1 \leq \frac{n}{2}$), the set of partners is increased by at least n new points. To prove the induction step, we shall use the following two definitions.

Definition 1: Let I be the image, and P a collection of points in the image. A *neighbor* of the set P is a point in $I - P$ that shares an edge or a vertex on the grid with a point in P .

Definition 2: In a grid of size n^2 , a set of k points is called *admissible* (and the grid itself is admissible) if:

$$\frac{1}{4}n^2 < k < \frac{3}{4}n^2 \quad \text{for } n > 2 \quad (2)$$

$$\frac{1}{4}n^2 \leq k < \frac{3}{4}n^2 \quad \text{for } n = 2$$

Comment: Let P be the set of the k original points, and all their partners for an operator of size M . When the operator increases in size to $M + 1$, all the neighbors of P become new partners of the k original points. To prove proposition 1 it will be sufficient, therefore, to show that the set P has at least n neighbors. This will be the goal of proposition 2.

Without loss of generality, we can assume that the set P is admissible. Since P contains the k original points, clearly $\frac{n^2}{4} < \|P\|$. If $\|P\| \geq \frac{3}{4}n^2$, then the k original points have at least $\frac{3}{4}n^2 - k$ partners. But since $k \leq \frac{n^2}{4} + 1$, and $\frac{n}{2} \geq M + 1$, it would follow that the k original points have at least $n \cdot m$ partners. Proposition 1 would therefore hold for the operator of size $M + 1$, and the induction step would be unnecessary.

Proposition 2: In a grid of n^2 points, any admissible set of K points has at least n neighbors.

Proof: By induction on n , the size of the grid. For $n = 2$ the proposition holds, since $K = 1$ or $K = 2$, and in either case there are at least 2 neighbors. We next assume that proposition 2 holds for $n = N$, and prove it for $n = N + 1$.

The general structure of the proof will be as follows. For convenience of reference, the K points will be called the black points, and the remaining points white. In these terms, we wish to show that in a grid of size $(N + 1)^2$ any set of K points, $\frac{1}{4}(N + 1)^2 < K < \frac{3}{4}(N + 1)^2$, has at least $N + 1$ white neighbors. To use the induction hypothesis we shall prove the following claim:

Proposition 3: It is possible to remove a row and a column from the grid in such a way, that the resulting grid of size N^2 will be admissible.

From the induction hypothesis, in the reduced grid the set of black points has at least N neighbors. The proof will be concluded by showing that when the missing row and column are re-introduced, the number of neighbors of the black set is increased by at least one.

Proposition 3.1: From an admissible grid of size $(N + 1)^2$ it is possible to remove a row and a column, so that in the remaining grid of size N^2 the number of black points will be less than $\frac{3}{4}N^2$.

Proof: Let α_{N+1} denote the fraction of black points in the original grid. That is, the number K of black points is equal to $\alpha_{N+1} \cdot (N + 1)^2$. Similarly, α_N is the fraction of black points in the reduced grid. In these terms, we assume in proposition 3.1 that $\frac{1}{4} < \alpha_{N+1} < \frac{3}{4}$, and wish to establish that $\alpha_N < \frac{3}{4}$.

Let B_{ij} denote the total number of black points in row i and column j . \bar{B}_{ij} is the (number of black points in row i) + (the number of black points in row j). The difference between B_{ij} and \bar{B}_{ij} is that if point (i,j) is black, it is counted twice in \bar{B}_{ij} but only once in B_{ij} . Let $B = \sum_{i,j=1}^{N+1} B_{ij}$, $\bar{B} = \sum_{i,j=1}^{N+1} \bar{B}_{ij}$.

There are $\alpha_{N+1}(N + 1)^2$ black points in the grid, and each one is counted $2(N + 1)$ times in \bar{B} , therefore:

$$\bar{B} = 2(N + 1) \cdot \alpha_{N+1}(N + 1)^2 \quad (3)$$

In \bar{B} each black point is counted one additional time, compared with B , therefore:

$$B = \bar{B} - \alpha_{N+1}(N + 1)^2 \quad (4)$$

There are $(N + 1)^2$ terms B_{ij} contributing to the sum B , therefore the average contribution is:

$$\frac{B}{(N + 1)^2} = 2\alpha_{N+1}(N + 1) - \alpha_{N+1} \quad (5)$$

There must exist a term B_{ij} that contributes at least the average contribution. Proposition 3.1 will therefore be satisfied if:

$$\alpha_{N+1}(N + 1)^2 - (2\alpha_{N+1}(N + 1) - \alpha_{N+1}) < \frac{3}{4}N^2 \quad (6)$$

If this inequality does not hold, then:

$$\alpha_{N+1}(N+1)^2 - (2\alpha_{N+1}(N+1) - \alpha_{N+1}) \geq \frac{3}{4}N^2 \quad (7)$$

But this implies $\alpha_{N+1} \geq \frac{3}{4}$, which is a contradiction, since $\frac{1}{4} < \alpha_{N+1} < \frac{3}{4}$. ■

Proposition 3.2: In the conditions of proposition 3.1 it is possible to remove a row and a column so that in the reduced grid $\alpha_N > \frac{1}{4}$.

The proof will be omitted, since it is entirely analogous to the proof of proposition 3.1.

Comment: The row and column chosen in propositions 3.1 and 3.2 are not necessarily the same. If, in proposition 3.1, $\alpha_N > \frac{1}{4}$, or if, in proposition 3.2, $\alpha_N < \frac{3}{4}$, then proposition 3 is established. Proposition 3 may still fail, however, to be satisfied if in proposition 3.1 $\alpha_N \leq \frac{1}{4}$ and in proposition 3.2 $\alpha_N \geq \frac{3}{4}$. It is required therefore to show that in this case we can still remove a row and a column such that in the reduced grid the set of black points is admissible.

Let the row and column chosen in proposition 3.1 be row i and column j . The row and column chosen in proposition 3.2 will be denoted row k and column l . Let us now remove from the grid row i and column l . Since the definition of an admissible set is slightly different for $N = 2$, we shall start with the case $N \geq 3$: In this case, proposition 3 would fail to be satisfied if either:

$$\alpha_{N+1}(N+1)^2 - B_{il} \leq \frac{1}{4}N^2 \quad (8)$$

or:

$$\alpha_{N+1}(N+1)^2 - B_{il} \geq \frac{3}{4}N^2 \quad (9)$$

Let us examine the first case. We may assume that

$$\alpha_{N+1}(N+1)^2 - B_{kl} \geq \frac{3}{4}N^2 \quad (10)$$

(see the comment above). From (8) and (10) we conclude that:

$$B_{il} - B_{kl} \geq \frac{1}{2}N^2 \quad (11)$$

$B_{il} - B_{kl}$ is the difference in the number of black points between rows i and k , after removing the l^{th} column, hence $B_{il} - B_{kl} \leq N$. This, together with (11), implies $N \leq 2$, in contradiction with the assumption that $N \geq 3$. We conclude that for $N \geq 3$ (8) is impossible, and the inequality $\alpha_{N+1}(N+1)^2 - B_{il} > \frac{1}{4}N^2$ must hold. To include the case $N = 2$, a similar argument establishes that $\alpha_{N+1}(N+1)^2 - B_{il} \geq \frac{1}{4}N^2$ must hold for $N \geq 2$.

The case in (9) can be handled analogously, yielding

$$B_{ij} - B_{il} > \frac{1}{2}N^2 \quad (12)$$

which again cannot hold for $N \geq 2$.

In conclusion, the fraction α_N satisfies $\frac{1}{4} < \alpha_N < \frac{3}{4}$ for $N \geq 3$, and $\frac{1}{4} \leq \alpha_N < \frac{3}{4}$ for $N = 2$. The set of black points in the reduced grid is therefore admissible. ■

Proposition 3 has established that a row and a column can be removed from the original grid of size $(N+1)^2$, resulting in an admissible grid of size N . From the induction hypothesis, in the reduced grid the set of black points has at least N neighbors. To show that the set of neighbors in the original grid is larger (by at least one), we shall use the following claim:

Proposition 4: proposition 3 can be satisfied by removing a row and a column that are not entirely black.

Proof: It is sufficient to show that proposition 3.1 can be satisfied with a row and a column that contain at least one white point. We can assume without loss of generality that the original grid contains the maximal possible number of black points, since if proposition 3.1 can be satisfied (with a row and a column that are not entirely black) for a maximal number of black points, it can also be satisfied for grids with fewer black points. For $N+1 = 2m$ (an even number), the maximal number of black points is $\frac{3}{4}(N+1)^2 - 1$. For

$N + 1 = 2m + 1$, the maximal number is $\frac{3}{4}(N + 1)^2 - \frac{3}{4}$. We shall examine the latter case (the first can be established analogously).

Let r be the number of completely black rows in the original grid, c the number of completely black columns. We can assume that $r \geq 1, c \geq 1$, since otherwise proposition 4 follows directly from proposition 3. Since $\alpha_{N+1} < \frac{3}{4}$, either $c < \frac{N+1}{2}$ or $r < \frac{N+1}{2}$ must hold. Without loss of generality we shall assume that $c < \frac{N+1}{2}$.

The r black rows and c black columns contribute a total of

$$(r + c)(N + 1) - rc \quad (13)$$

black points. The number of black points in the remaining rows and columns is therefore:

$$\frac{3}{4}(N + 1)^2 - \frac{3}{4} - (r + c)(N + 1) + rc \quad (14)$$

This number of black points is divided among $N + 1 - c$ columns, hence the average per column (denoted by A) is:

$$A = \frac{\frac{3}{4}(N + 1)^2 - \frac{3}{4} - (r + c)(N + 1) + rc}{N + 1 - c} \quad (15)$$

There must exist a column which contains at least $r + A$ black points. By removing this column and one of the black rows we remove at least:

$$(N + 1) + r + A - 1 \quad (16)$$

black points. It therefore remains to establish that:

$$\frac{3}{4}(N + 1)^2 - \frac{3}{4} - (N + r + A) < \frac{3}{4}N^2 \quad (17)$$

Inequality (17) reduces to:

$$\frac{N^2}{4} + N - \frac{Nc}{2} - c > 0 \quad (18)$$

For $c < \frac{N+1}{2}$ inequality (18) holds, therefore proposition 4 is established. ■

Corollary: using the same argument, proposition 3.2 can be established with a row and a column that are not entirely white. Combining the two claims, it is straightforward to establish that proposition 3 can be satisfied by a row and a column that are neither entirely black nor entirely white.

Proposition 5: The set of black points in the original grid has at least $N + 1$ neighbors.

Proof: From propositions 3 and 4, we can remove a row R and a column C (neither entirely black nor entirely white) from the original grid, and obtain an admissible grid. From the induction hypothesis, the set of black points in the reduced grid has at least N neighbors. We have to show that by re-introducing R and C the number of neighbors increases by at least one.

Since R and C contain both black and white points, they contain at least one "internal neighbor", i.e., a white point that borders a black point, where both points are in $R \cup C$. We shall show that without the internal neighbor the number of neighbors in the original grid is at least N , hence the total number of neighbors is at least $N + 1$, as required.

If a given point p is a neighbor of the black set in the reduced grid, it may or may not remain a neighbor when the missing row and column are re-introduced. Let us examine first the case where p is not a neighbor of the black set in the original grid. In this case p must be surrounded in the original grid by white points (see Figure 2). In this configuration, point q in R becomes a new neighbor, since at least one of the points a, b, c must be black. Note that q cannot be an internal neighbor. (Comment: in Figure 2, point p borders the re-introduced column R . The argument is unchanged when it borders C instead of R . The case where p borders both R and C requires a slight extension of the argument, which will not be elaborated.)

We conclude that there exists a one-to-one mapping from the neighbors in the reduced grid into the neighbors in the original grid, and that this mapping does not include the internal neighbor. The number of neighbors in the grid of size $N + 1$ is therefore at least $N + 1$. This establishes the induction step in proposition 2, and concludes the proof of propositions 2 and 1. ■




Figure 2: After column **R** has been re-introduced, point p is no longer a neighbor of the black set. All the squares bordering p must therefore be white. Prior to the re-introduction of column **R**, p was a neighbor of the black set, hence at least one of the points a, b, c , must be black. Point q is therefore a neighbor of the black set.

4. Reducing the size of the intermediate storage

The lower bound established in the preceding section implies that the intermediate storage may play a dominant role in determining the size of the system. The intermediate storage can, however, be reduced by scanning the image more than once. Suppose, for example, that s rows are scanned in parallel. The first scan-path scans rows 1 to s , the second scans rows $s + 1$ to $2s$, etc. Each image point is scanned in this scheme s times. Using the serpentine delay line of Section 2, it is possible now to connect s lines directly to the two-dimensional operator M . The intermediate storage S will then have only $(m - s)$ rows. This scheme gives a simple (though not optimal, in terms of the storage size) method of reducing the size of the intermediate storage to $n \cdot (m - s)$, where s is the number of times each point is scanned. In particular, by having $s = m$ the intermediate storage becomes unnecessary.

References

- Hall, J. E. & Awtrey, J. D. Real-time image enhancement using 3×3 pixel neighborhood operator functions. *Paper presented at the SPIE Symposium, San Diego, California, August 1979.*
- Nudd, G. R., Nygaard, P. A., Fouse, S. D., & Nussmeier. Implementation of advanced real-time image understanding algorithms. in: L. S. Baumann (Ed.) *Proceedings of Image Understanding Workshop, Menlo Park, California, pp. 151-158 April 1979.*

Acknowledgement: I thank M. Brady and E. Hildreth for their helpful comments.

